

4 Literatur

- ANDRÁSSY I. (1984): Klasse Nematoda (Ordnungen Monhysterida, Desmoscolecida, Araeolaimida, Chromadorida, Rhabditida). Gustav Fischer Verlag, Stuttgart
- BONGERS T. (1988): De Nematoden van Nederland. Natuurhistorische Bibliotheek van de KNNV, Utrecht
- DEBÜS R., HUND K. (1995): Entwicklung analytischer Methoden zur Erfassung biologisch relevanter Belastungen von Böden. Statusseminar zum Förderschwerpunkt „Ökotoxikologie“ des BMBF 2/95: 215–225
- DONKIN S.G., DUSENBERY D.B. (1993): A soil toxicity test using the nematode *Caenorhabditis elegans* and an effective method of recovery. Archives of Environmental Contamination and Toxicology 25: 145–151
- DONKIN S.G., DUSENBERY D.B. (1994): Using the *Caenorhabditis elegans* soil toxicity test to identify factors affecting toxicity of four metal ions in intact soil. Water Air & Soil Pollution 78: 359–373
- FREY F. (1976): Untersuchungen über die Wirkungen von im Obstbau verwendeten Herbiziden auf den Testnematoden *Acrobeloides buteschlii* (de Man, 1884) Steiner und Buhrer, 1933. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 83: 434–441
- GILBERTSON M. (1989): Effects on fish and wildlife populations. In: R. KIMBROUGH & A.A. JENSEN (eds.) *Halogenated biphenyls, terphenyls, naphthalenes, dibenzodioxins and related products*. Elsevier Science Publ., pp. 103–127
- GUVEN K., DUCE J.A., DE POMERAI D.I. (1994): Evaluation of a stress-inducible transgenic nematode strain for a rapid aquatic toxicity testing. Aquatic Toxicology 29: 119–137
- HAIGHT M., MUDRY T., PASTERNAK J. (1982): Toxicity of seven heavy metals on *Panagrellus silusiae*: the efficacy of the free-living nematode as an *in vivo* toxicological bioassay. Nematologica 28: 1–11
- HEIMBACH F. (1985): Comparison of laboratory methods, using *Eisenia foetida* and *Lumbricus terrestris*, for the assessment of the hazard of chemicals to earthworms. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 92: 156
- HOY J.B. (1990): Long-term effects of lindane on the oribatid community in a pine plantation. Environmental Entomology 19: 898–900
- KAMMENGA J.E., VAN GESTEL C.A.M., BAKKER J. (1994): Patterns of sensitivity to cadmium and pentachlorophenol among nematode species from different taxonomic and ecological groups. Archives of Environmental Contamination and Toxicology 27: 88–94
- KÖRDEL W., WAHLE U. (1995): Methoden zur Erfassung biologisch relevanter Belastungen von Böden – Extraktion und chemische Analytik biologisch relevanter Belastungen. Abschlußbericht, Förder- schwerpunkt Ökotoxikologie des BMFT, eingereicht.
- MACEK K.J., BUXTON K.S., DERR S.K., DEAN J.W., SAUTER S. (1976): Chronic toxicity of lindane to selected aquatic invertebrates and fish. Washington D.C., US-EPA Ecological Research Series EPA-600/3-76-046
- MAYER F.L., MEHRLE P.M., SANDERS H.O. (1977): Residue dynamics and biological effects of polychlorinated biphenyls in aquatic organisms. Archives of Environmental Contamination and Toxicology 5: 501–511
- NICHOLAS W.L. (1975): The biology of free-living nematodes. Clarendon Press, Oxford
- PAGANO G., CIPOLLARO M., CORSALE G., ESPOSITO A., RAGUCCI E., GIORDANO G.G., TRIEFF N.M. (1985): Comparative toxicities of chlorinated biphenyls on sea urchin egg fertilisation and embryogenesis. Marine Environmental Research 17: 240–244
- PARMELEE R.W., WENTSEL R.S., PHILLIPS C.T., SIMINI M., CHECKAI R.T. (1994): Soil microcosm for testing the effects of chemical pollutants on soil fauna communities and trophic structure. Environmental Toxicology and Chemistry 12: 1477–1486
- POPHAM J.D., WEBSTER J.M. (1982): Ultrastructural changes in *Caenorhabditis elegans* (Nematoda) caused by toxic levels of mercury and silver. Ecotoxicology and Environmental Safety 6: 183–189
- SAFE S. (1992): Toxicology, structure-function relationship, and human and environmental health impacts of polychlorinated biphenyls: progress and problems. Environmental Health Perspectives 100: 259–268
- SAMOILOFF M.R. (1990): The nematode toxicity assay using *Panagrellus redivivus*. Biological Assessment 5: 309–318
- SLOOFF W., CANTON J.H., HERMENS J.L.M. (1983): Comparison of the susceptibility of 22 freshwater species to 15 chemical compounds. I. (Sub)acute toxicity tests. Aquatic Toxicology 4: 113–128
- SMITH V.L., JOHNSTON P.A. (1992): Differential haemotoxic effect of PCB congeners in the common shrimp, *Crangon crangon*. Comparative Biochemistry and Physiology 101 C: 641–649
- STALLING D.L., MAYER F.L. jr. (1972): Toxicities of PCBs to fish and environmental residues. Environmental Health Perspectives 1: 159–164
- STRINGHAM E.G., CANDIDO E.P.M. (1994): Transgenic hsp16-lacZ strains of the soil nematode *Caenorhabditis elegans* as biological monitors on environmental stress. Environmental Toxicology and Chemistry 13: 1211–1220
- SUDHAUS W. (1976): Vergleichende Untersuchungen zur Phylogenie, Systematik, Ökologie, Biologie und Ethologie der Rhabditidae (Nematoda). Zoologica 43: 1–229
- THURSTON R.V., GILFOIL T.A., MEYN E.L., ZAJDEL R.K., AOKI T.I., VEITH G.D. (1985): Comparative toxicity of ten organic chemicals to ten common aquatic species. Water Research 19: 1145–1155
- VILLE P., ROCH P., COOPER E.L., MASSON P., NARBONNE J.F. (1995): PCBs increase molecular-related activities (lysozyme, antibacterial, hemolysis, proteases) but inhibit macrophage-related functions (phagocytosis, wound healing) in earthworms. Journal of Invertebrate Pathology 65: 217–224
- WASILEWSKA L. (1989): Impact of human activities on nematode communities in terrestrial ecosystems. In: B. CLARHOLM & L. BERGSTROM (eds.) *Ecology of Arable Land – Perspectives and Challenges*. Kluwer Acad. Publ., Dordrecht, pp. 123–132
- WHO (1991): Environmental Health Criteria 124: Lindane. World Health Organization, Geneva
- WILLIAMS P.L., DUSENBERY D.B. (1988): Using the nematode *Caenorhabditis elegans* to predict mammalian acute lethality to metallic salts. Toxicology and Industrial Health 4: 469–478
- WILLIAMS P.L., DUSENBERY D.B. (1990): Aquatic toxicity testing using the nematode, *Caenorhabditis elegans*. Environmental Toxicology and Chemistry 9: 1285–1290
- WILMS W. (1992): Prüfung der Auswirkungen von Pflanzenschutzmitteln auf räuberische Nematoden. Nachrichtenblatt des Deutschen Pflanzenschutzdienstes 44: 25–29
- YEATES G.W., BONGERS T., DE GOEDE R.G.M., FRECKMAN D.W., GEORGIEVA S.S. (1993): Feeding habits in nematode families and genera – an outline for soil ecologists. Journal of Nematology 25: 315–331

Eingegangen: 20. 12. 1995
Akzeptiert: 18. 09. 1996

Nematoden in der ökotoxikologischen Forschung – Plädoyer für eine vernachlässigte, jedoch sehr aussagekräftige Tiergruppe

UWSF 7 (2) 74–83 (1995)

W. TRAUNSPURGER, CHR. STEINBERG, T. BONGERS

Die ökologische Bedeutung freilebender Nematoden (Fadenwürmer) wird dargestellt und ein ataxonomischer sowie taxonomischer Bioindikationsansatz diskutiert.