

RESEARCH

Open Access



# Study on the influence mechanism of underground mineral element Fe(II) on Cr(VI) transformation under subsurface and groundwater interaction zones

Yiru Zheng<sup>1</sup>, Shuguang Liu<sup>1,2</sup>, Chaomeng Dai<sup>1\*</sup>, Yanping Duan<sup>3\*</sup>, Aleksei Nikolavich Makhinov<sup>4</sup>, Leong Kah Hon<sup>5</sup> and José Tavares Araruna Júnior<sup>6</sup>

## Abstract

**Background:** With the acceleration of urbanization, heavy metal contamination in subsurface water and groundwater is becoming more and more serious, and the interaction between surface water and underground water is an important factor that cannot be ignored. Based on the change of oxygen content in surface water and groundwater environment, this research mainly studied the influencing factors and mechanism of underground mineral element Fe(II) on the morphological transformation of heavy metal Cr(VI) under the condition of surface–groundwater interaction, and simulates the interaction process of subsurface–groundwater by changing the dissolved oxygen content in the redox reaction system of dissolved Fe(II) and Cr(VI). The influence mechanism of pH, Fe(II):Cr(VI), groundwater hardness ion and humic acid on the morphological transformation of Cr(VI) was investigated.

**Results:** The results showed that the content of DO did affect the conversion of Cr(VI) by Fe(II). pH had a great influence on the redox reaction: while pH = 3–5, the reaction rate of Fe(II) and Cr(VI) and the removal rate of Cr(VI) decreased with the increase of pH; while pH = 6–8, the reaction rate and the removal of Cr(VI) increased with the increase of pH. When Fe(II):Cr(VI) = 3:1, the reaction of Fe(II) and Cr(VI) is affected by DO; when Fe(II):Cr(VI) is 1:1, 2:1, 4:1 and 5:1, DO has no obvious effect on the reaction. Groundwater hardness ions can greatly promote the redox reaction. When humic acid exists in the reaction system, the Fe(III) formed by the oxidation of Fe(II) by Cr(VI) will once again be converted into soluble Fe(II), the reduction of humic acid further promotes the redox reaction between Fe(II) and Cr(VI), and promotes the conversion of Cr(VI) by Fe(II) more effectively. Moreover, the higher the concentration of humic acid, the higher the removal rate of Cr(VI).

**Conclusions:** The research results have practical guiding significance for groundwater contamination prediction, pollution control and remediation, integrated water resources management and the construction of sponge city in China.

**Keywords:** Subsurface–groundwater interaction, Redox reaction, Hexavalent chromium, Ferrous iron, Dissolved oxygen

## Background

Hexavalent chromium is one of the most widely distributed elements in nature, which is used in various industrial processes and industrial products, such as mining smelting, chemical industry, paint pigment, pharmacy, light industry textile, leather tanning agent and so on.

\*Correspondence: chaomeng.dai@tongji.edu.cn; duanyanping@shnu.edu.cn

<sup>1</sup> Department of Hydraulic Engineering, Tongji University, Shanghai, China

<sup>3</sup> School of Environmental and Geographical Sciences, Shanghai Normal University, Shanghai, China

Full list of author information is available at the end of the article

Then as industrial waste, hexavalent chromium is discharged into the water environment, which has a serious impact on the water ecological environment and even the human body [1, 2]. Nevertheless, the valence of chromium mainly exists in the form of +3 and +6 in the water environment [3, 4]. Among them, Cr(III) is not only less toxic, stable, not easy to migrate, but also a necessary trace element for human body. Compared with Cr(III), Cr(VI) has stronger mobility and toxicity [4], and is a strong oxidant and potential carcinogen. After entering the surface water environment, heavy metal pollution in surface runoff is formed, and it is constantly moving down, which threatens the groundwater environment [4]. In the process of natural hydrological cycle, the interaction between surface water and groundwater is a common phenomenon [5], and plays an important role [6]. Almost all surface water bodies in nature interact with groundwater. Not only the exchange of water quantity, but also the transport and transformation of solute and pollutants [7], this interaction will have a significant impact on the migration and transformation of chromium, a heavy metal pollutant. Therefore, the interaction between surface and groundwater is an important factor that cannot be ignored in the study of the transport and transformation of heavy metal pollutants. At present, the research on the effect of groundwater interaction on the water quality at home and abroad mainly focuses on the change of the water chemistry environment on the surface-ground water interaction, thus affecting the complex biogeochemical process of the interaction belt, and changing the pH, Eh, DO flowing through the water, and the form of matter and the transformation of migration [8–10]. Some scholars have found that the surface-ground water interaction process The transport and conversion of medium heavy metal pollutants are subjected to convection, diffusion, dispersion, adsorption and desorption, and some chemical reactions, such as oxidation factors such as reduction reaction, hydrolysis and biotransformation [11]. Generally speaking, it can be summarized as the following three points; (1) the direction of the interaction between subsurface water and groundwater with water and contaminants [12–15]. (2) The transport mechanism of contaminants, including convection, diffusion, dispersion, dispersion and adsorption, and desorption. (3) Biological/chemical reactions, including irreversible reactions such as abiotic reactions, biodegradation and radioactive decay, and reversible reactions such as precipitation dissolution, adsorption, redox and ion exchange [12, 16]. Eh-sensitive mineral elements such as Fe, S, Mn and organic matter are found in natural underground environment [17], so, normally, the underground environment medium is reductive. When

the subsurface water interacts with the groundwater, the largest change is contact with the oxidizing substance (usually  $O_2$ ), which makes the Eh significantly higher [6, 18, 19]. In this regard, Singh et al. [20] studied the effects of different pH on the redox reaction of Fe (II) with heavy metal Cr(VI) in water under anoxic and aerobic conditions. The results showed that the main reason why Cr(VI) could not be completely reduced in the presence of oxygen was that Fe(II) was oxidized to Fe(III). Stumm et al. [20] have found that the redox cycle between Fe(II)/Fe(III) can play the role of the electronic shuttle in a large amount of biological and non-biological reaction, and further influence the migration and transformation of heavy metal pollutants in the underground environment. Eh, which can obviously change the underground medium environment in the interaction process between natural surface water and groundwater, makes the reductive substances such as Fe(II), which widely exist in the underground environment, easily redox cycle with heavy metal pollutants Cr(VI) with the change of Eh, which will further affect the morphological transformation of Cr(VI) into groundwater environment. At present, the morphological transformation of heavy metal pollutants under the interaction of subsurface groundwater has not been studied in domestic and foreign studies.

Therefore, the influence factors and mechanism of Fe(II) on the transformation of heavy metal Cr(VI) in the underground mineral element Fe(II) under the condition of subsurface-underground water interaction are mainly studied. The highlight point of this research lies in simulating the subsurface-groundwater interaction process by altering the dissolved oxygen content in the solubility Fe(II) and Cr(VI) redox reaction system. On this basis, static experiments were used to explore the mechanism of the effect of initial solution pH, compared with Fe(II):Cr(VI), groundwater hardness ions and humic acid on the redox reaction of Fe(II) in Cr(VI).

## Materials and methods

### Materials

All chemicals used in this experiment were of analytical grade, and all stock solutions were prepared with deionized (DI) water from a Milli-Q water system. The Cr(VI) stock solutions were prepared by dissolving a weighted amount of dried  $K_2CrO_4$  (Sinopharm, AR grade) in DI water. Fe(II) stock solutions were prepared in an anaerobic chamber just before use by dissolving a weighed amount of dried  $Fe(NH_4)_2(SO_4)_2$  (Sinopharm, AR grade) in DI water. The initial reaction temperature is about  $(20 \pm 1) ^\circ C$ . pH adjustment of solutions was made by 0.1 M  $HNO_3$  and 0.1 M NaOH.

**Table 1 Effect of dissolved oxygen experiment**

Case	Experiment conditions (nitrogen blowing) (min)	$C_{Fe(II)}$ (mg/L)	$C_{Cr(VI)}$ (mg/L)	DO (mg/L)
1	0	20	20	Hyperoxia (5.95)
2	15	20	20	Hypoxia (3.98)
3	30	20	20	Hypoxia (2.21)
4	60	20	20	Hypoxia (1.20)
5	120	20	20	Anoxic (0.11)

**Table 2 Effect of initial pH experiment**

Case	Experiment conditions	$C_{Fe(II)}$ (mg/L)	$C_{Cr(VI)}$ (mg/L)	DO
1	pH=3	20	20	Hyperoxia
2		20	20	Anoxic
3	pH=4	20	20	Hyperoxia
4		20	20	Anoxic
5	pH=5	20	20	Hyperoxia
6		20	20	Anoxic
7	pH=6	20	20	Hyperoxia
8		20	20	Anoxic
9	pH=7	20	20	Hyperoxia
10		20	20	Anoxic
11	pH=8	20	20	Hyperoxia
12		20	20	Anoxic

### Batch experiments

This experiment was carried out in a batch experiment, and all the reactions were carried out in a three-necked flask [21]. It can be divided into five parts: (1) the dissolved oxygen content in the reaction system was controlled by changing the blowing time of nitrogen. The blowing time was 0 min, 15 min, 30 min, 60 min and 120 min, respectively. The corresponding DO concentration was measured by dissolved oxygen meter to simulate the interaction process of surface-groundwater (Table 1). (2) The pH gradient of the initial solution was 3.0, 4.0, 5.0, 6.0, 7.0 and 8.0. The gradient was adjusted by 0.1 M concentrated nitric acid and 0.1 M sodium hydroxide solution and determined by pH-3E acidometer (Table 2). (3) In this part, the effect of initial reactant concentration ratio on the morphological transformation of Cr(VI) was studied. Fe(II):Cr(VI) was set to 1:1, 2:1, 3:1, 4:1 and 5:1. To meet the actual pH environment of groundwater, pH was controlled to 6 (Table 3). (4) The hardness ions in groundwater, namely calcium, magnesium and bicarbonate ions, were investigated. The effect of Fe(II) on the morphological transformation of Cr(VI) was studied by changing the concentration of these ions.  $CaCl_2$ ,  $MgCl_2$  and  $NaHCO_3$  were used as reaction reagent in the

**Table 3 Effect of initial reactant concentration ratio Fe(II):Cr(VI)**

Case	Experiment conditions	$C_{Fe(II)}$ (mg/L)	$C_{Cr(VI)}$ (mg/L)	DO
1	Fe(II):Cr(VI) = 1:1	20	20	Hyperoxia
2		20	20	Anoxic
3	Fe(II):Cr(VI) = 2:1	45	20	Hyperoxia
4		45	20	Anoxic
5	Fe(II):Cr(VI) = 3:1	64	20	Hyperoxia
6		64	20	Anoxic
7	Fe(II):Cr(VI) = 4:1	86	20	Hyperoxia
8		86	20	Anoxic
9	Fe(II):Cr(VI) = 5:1	106	20	Hyperoxia
10		106	20	Anoxic

experiment (Table 4). (5) This part mainly investigated the effect of natural organic matter represented by humus on the morphological transformation of Cr(VI), which was the most widely distributed natural organic matter in ecological environment. As the main component, humic acid was used in the experiment [22]. The concentration gradient of humic acid was set to 0 mg/L, 15 mg/L and 40 mg/L, respectively, as well as the blank control group experiment (Table 5).

### Analytical methods

At each sampling time, the 10 mL sample was taken from the opening at the upper end of the device with a syringe, and added 2,2'-bipyridine of 3 nM to block the continuation of the redox reaction between Fe(II) and Cr(VI) in the sample immediately (Tong 2015), and then the sample was filtered with a microporous filter of 0.22  $\mu$ m. The Cr(VI) content in the reaction system was determined according to the determination of hexavalent chromium in water quality by diphenylcarbazide spectrophotometry (GB7487-87). The filtered sample was placed in 50-mL colorimetric tube. The sulfuric acid solution of 2.5 mL (1+7) and the diphenylcarbamide dihydrazide acetone solution of 2.5 mL were added successively, and then diluted to 50 mL with water. After shaking for 5–10 min, the absorbance was

**Table 4 Effect of groundwater hardness experiment**

Case	Experiment conditions	C <sub>Fe(II)</sub> (mg/L)	C <sub>Cr(VI)</sub> (mg/L)	C <sub>Ca<sup>2+</sup></sub> (mg/L)	C <sub>Mg<sup>2+</sup></sub> (mg/L)	C <sub>HCO<sub>3</sub><sup>-</sup></sub> (mg/L)	DO
1	Blank	20	20	0	0	0	Hyperoxia
2		20	20	0	0	0	Anoxic
3	Only Ca <sup>2+</sup>	20	20	40	0	0	Hyperoxia
4		20	20	40	0	0	Anoxic
5	Only Mg <sup>2+</sup>	20	20	0	40	0	Hyperoxia
6		20	20	0	40	0	Anoxic
7	Only HCO <sub>3</sub> <sup>-</sup>	20	20	0	0	60	Hyperoxia
8		20	20	0	0	60	Anoxic
9	Ca <sup>2+</sup> , HCO <sub>3</sub> <sup>-</sup>	20	20	40	0	60	Hyperoxia
10		20	20	40	0	60	Anoxic

**Table 5 Effect of humic acid experiment**

Case	Experiment conditions	C <sub>Fe(II)</sub> (mg/L)	C <sub>Fe(III)</sub> (mg/L)	C <sub>Cr(VI)</sub> (mg/L)	C <sub>Cr(III)</sub> (mg/L)	HA (mg/L)	DO
1	HA + Fe(II)	20	0	0	0	40	Hyperoxia
2		20	0	0	0	40	Anoxic
3	HA + Fe(III)	0	12	0	0	40	Hyperoxia
4		0	12	0	0	40	Anoxic
5	HA + Cr(III)	0	0	0	12	40	Hyperoxia
6		0	0	0	12	40	Anoxic
7	HA + Cr(VI)	0	0	20	0	40	Hyperoxia
8		0	0	20	0	40	Anoxic
9	HA = 0 mg/L	20	0	20	0	0	Hyperoxia
10		20	0	20	0	0	Anoxic
11	HA = 15 mg/L	20	0	20	0	15	Hyperoxia
12		20	0	20	0	15	Anoxic
13	HA = 40 mg/L	20	0	20	0	40	Hyperoxia
14		20	0	20	0	40	Anoxic

determined with pure water as reference at the wavelength of 542 nm, and the concentration of Cr(VI) was calculated compared with the standard line. Fe(II) was measured by 1,10-phenanthroline spectrophotometry (wavelength 510 nm).

#### Calculation methods

The redox reaction rate of Fe(II) and Cr(VI) can be represented by a pseudo-first-order reaction model [23].

$$\frac{-d[\text{Cr(VI)}]}{dt} = k[\text{Cr(VI)}][\text{Fe(II)}]. \quad (1)$$

In this reaction, taking Fe(II):Cr(VI)=1:1 as an example, the reaction rate equation can be rewritten into


$$\frac{1}{[\text{Cr(VI)}]_t} = \frac{1}{[\text{Cr(VI)}]_0} + kt \quad (2)$$

where  $k$  is the apparent rate constant,  $[\text{Cr(VI)}]_0$  the initial concentration of Cr(VI) and  $[\text{Cr(VI)}]_t$  the concentration corresponding to reaction time of Cr(VI).

## Results and discussion

### Effects of dissolved oxygen (DO)

The dissolved oxygen content in the reaction system was controlled by changing the nitrogen blowing time. According to the amount of dissolved oxygen concentration, the change of dissolved oxygen content was divided into three types of hyperoxia, hypoxia and anoxic, which are defined as follows: (1) hyperoxia: the nitrogen blowing time was 0 min, DO=5.95 mg/L; (2) hypoxia: the nitrogen blowing time was 15 min, 30 min, 60 min, and the corresponding DO concentrations were 3.98 mg/L, 2.21 mg/L, 1.20 mg/L, respectively; (3) anoxic: the nitrogen blowing time was 120 min, DO=0.11 mg/L. The mechanism of the influence of underground mineral





element Fe(II) on the morphological transformation of heavy metal pollutants Cr(VI) was simulated during the interaction of surface-groundwater. The initial concentration of Cr(VI) and Fe(II) was 20 mg/L, the total react time was 160 min, as shown in Fig. 1.

It can be seen from the figure that the concentrations of Cr(VI) and Fe(II) in the reaction system decreased rapidly in the pre-5 min before the reaction began, indicating that the reaction occurred in an instant [24]. Under the conditions of high oxygen, low oxygen and oxygen, the time for the complete reaction equilibrium was 120 min, 60 min and 20 min, respectively, but the concentration of Cr(VI) in the late stage of oxygen-free condition was gradually increased, and the release phenomenon was observed. The reason was that under the condition of anoxic as well as Fe(II):Cr(VI) = 1:1, the amount of Fe(II) is insufficient, and the oxidation of Cr(VI) in the early stage of the reaction can be carried out rapidly according to the equilibrium principle of the redox reaction. The reaction is carried out in the reverse direction at the later stage. The concentration of Cr(VI) in the later stage reaction system has a tendency to increase. However, it was found that compared with anoxic, Fe(II) was more beneficial to Cr(VI) fixation under oxic conditions. Because Fig. 2 shows that in the absence of oxygen, the value of pH in the reaction system is lower than that in aerobic condition, which indicates that the pH under aerobic condition is relatively high, which is more beneficial to the formation of Cr(III), that is to say, it is more beneficial for Fe(II) to fix Cr(VI).

#### Effects of initial pH

The redox reaction between Fe(II) and Cr(VI) can be represented by formula (3):




It is well known that pH is an important factor in this redox reaction. The morphology of Fe(II) and Cr(VI) in water are different under different pH conditions, which will affect the reaction process and reaction rate [25]. Buerge et al. [23] have studied the kinetics of the reaction between Fe(II) and Cr(VI) in the range of pH = 2–7.2 without oxygen, but the effects of strong alkalinity and aerobic conditions on the redox reaction of Fe(II) and Cr(VI) have not been studied. As a matter of fact, in the process of surface-groundwater interaction, the concentration of DO changes, so the pH gradients of 3.0, 4.0, 5.0, 6.0, 7.0 and 8.0 are set up, and the cases of hyperoxic (a) and anoxic (b) are discussed, respectively. (To express the change effect of DO more clearly, we only selected hyperoxic and anoxic to carry on the control experiment, as well as the following experiments.) The initial concentration of Cr(VI) and Fe(II) was 20 mg/L, the total react time was 160 min, as shown in Fig. 3.


From the view of pH, the effects of different pH gradients on the morphology of Cr(VI) and the rate of Fe(II) and Cr(VI) redox reaction were analyzed. From the calculation of reaction kinetics and the removal rate of Cr(VI) (Figs. 3, 4 and Table 6), the pH has a great effect on the reaction (in the case of hyperoxia): pH = 3–5, the reaction rate constant  $k$  decreased from 0.198 to 0.159, and the removal rate of Cr(VI) decreased from 65.33 to 55.57%. The lowest value was appeared at pH = 5; pH = 6–8, the reaction rate was increased from 0.167 to 0.238, and Cr(VI) removal rate increased from 59.35 to 71.62%. The above results indicate that the reaction rate of Fe(II) and Cr(VI) decreased with the increase of pH at pH = 3–5, and the reaction rate of Fe(II) and Cr(VI) increased with the increase of pH at pH = 6–8, as well as Cr(VI) removal rate, which are consistent with previous studies [23, 26–28].

The effects of Fe(II) and Cr(VI) reactions under different pH gradients were analyzed from the view of the





**Fig. 3** Variation diagram of Fe(II) and Cr(VI) concentrations under different pH conditions



**Fig. 4** Removal rate of Cr(VI) under different pH conditions

change content of DO. Dissolved oxygen will be present in the redox environmental systems and may serve to interfere in the reaction between Cr(VI) and Fe(II) by its own ability to oxidize Fe(II). It is known that Fe(II) oxidation by DO is primarily dependent on the pH and the DO concentration of an aqueous solution [29]. Our

**Table 6** Calculating apparent constant of Fe(II) and Cr(VI) under different pH conditions

Case	pH	C <sub>Cr(VI)</sub> (mg/L)	k (Ms <sup>-1</sup> )	t <sub>1/2</sub> (s)		
Hyperoxia	3	20	0.198	3.502		
	4		0.167	4.159		
	5		0.159	4.363		
	6		0.167	4.159		
	7		0.170	4.068		
	8		0.238	2.915		
	Anoxic		3	20	0.185	3.751
			4		0.189	3.661
5		0.153	4.522			
6		0.161	4.295			
7		0.188	3.683			
8		0.210	3.299			

experimental results reveal that the existence of oxygen does affect the redox reaction of Fe(II) with Cr(VI) under different pH conditions, which is consistent with the results of Stumm and Schlautman et al.

When pH=7 is high oxygen or no oxygen, the concentration of Fe(II) and Cr(VI) in the solution changes smoothly, which indicates that the reduction effect is better under neutral conditions. Figure 3a describes that, in the case of hyperoxia, the concentration of Cr(VI) in the reaction system with a pH of 3 to 5 fluctuates, but the overall presentation of the decrease is due to the fact that, in the case of strong acidity, the Fe(II) is not oxidized by oxygen. It is the cause that the thermodynamics of Fe(II) cations in acidic solution is unstable and tends to be oxidized by oxygen. From Table 7, it is known that  $O_2 + 4H^+ + 4e^- = 2H_2O$ ,  $E^0 = 1.229$  V, but in kinetics, the oxidation reaction is slow. When there is strong oxidant potassium dichromate in the reaction system,  $HCrO_4^- + 7H^+ + 3e^- = Cr^{3+} + 4H_2O$ ,  $E^0 = 1.35$  V > 1.229 V [30], so Fe(II) is superior to  $O_2$  and Cr(VI) in redox reaction in acidic environment.

In the system with pH=6–8, the concentration of Fe(II) decreased rapidly, compared with that in acidic condition, and the removal rate of Cr(VI) was also higher than that in acidic condition (Fig. 4). However, the removal rate of Cr(VI) under anaerobic conditions is higher than that under high oxygen environment, and in the case of hyperoxia, the concentration of Cr(VI) in the solution gradually increases with the continuous exchange of oxygen in the later stage of the reaction. There were two reasons for this: on the one hand, as can be seen from Table 7, the standard electric potential  $E^0 = 0.771$  V for  $Fe^{3+}/Fe^{2+}$  is kept constant within a certain pH range without the precipitation of the hydrated oxide. But when the pH of the hydrated

**Table 7 Standard electrode potential meter**


Acid condition	
(a) $\text{Fe}^{3+} + \text{e}^- = \text{Fe}^{2+}$	$E^0 = 0.771 \text{ V}$
(b) $\text{O}_2 + 4\text{H}^+ + 4\text{e}^- = 2\text{H}_2\text{O}$	$E^0 = 1.229 \text{ V}$
(c) $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- = 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	$E^0 = 1.33 \text{ V}$
(d) $\text{HCrO}_4^- + 7\text{H}^+ + 3\text{e}^- = \text{Cr}^{3+} + 4\text{H}_2\text{O}$	$E^0 = 1.35 \text{ V}$
Basic condition	
(a) $\text{Cr}(\text{OH})_3 + 3\text{e}^- = \text{Cr} + 3\text{OH}^-$	$E^0 = -1.48 \text{ V}$
(b) $\text{CrO}_2^- + 2\text{H}_2\text{O} + 3\text{e}^- = \text{Cr} + 4\text{OH}^-$	$E^0 = -1.2 \text{ V}$
(c) $\text{Fe}(\text{OH})_2 + 2\text{e}^- = \text{Fe} + 2\text{OH}^-$	$E^0 = -0.877 \text{ V}$
(d) $\text{Fe}(\text{OH})_3 + \text{e}^- = \text{Fe}(\text{OH})_2 + \text{OH}^-$	$E^0 = -0.56 \text{ V}$
(e) $\text{CrO}_4^{2-} + 4\text{H}_2\text{O} + 3\text{e}^- = \text{Cr}(\text{OH})_3 + 5\text{OH}^-$	$E^0 = -0.13 \text{ V}$
(f) $\text{O}_2 + \text{H}_2\text{O} + 4\text{e}^- = 4\text{OH}^-$	$E^0 = 0.401 \text{ V}$

oxide precipitation occurs, the half-reaction of the electric pair becomes  $\text{Fe}(\text{OH})_3 + \text{e}^- = \text{Fe}(\text{OH})_2 + \text{OH}^-$ ,  $E^0 = -0.56 \text{ V}$ . In the alkaline environment, Fe(III) is more likely to form hydroxide precipitate than Fe(II). At this time, the potential is changed obviously, so that the reduction of Fe(II) is suddenly increased, and the oxidation of Fe(III) is decreased. Under the equilibrium of chemical reaction, the reaction formula is more likely to be carried out in the positive reaction direction.


Therefore, the removal rate of Cr(VI) in alkaline environment is higher than that in acidic environment. It is also because in alkaline environment,  $\text{O}_2 + \text{H}_2\text{O} + 4\text{e}^- = 4\text{OH}^-$ ,  $E^0 = 0.401 \text{ V}$ ,  $\text{CrO}_4^{2-} + 4\text{H}_2\text{O} + 3\text{e}^- = \text{Cr}(\text{OH})_3 + 5\text{OH}^-$ ,  $E^0 = -0.13 \text{ V} < 0.401 \text{ V}$ , indicating that the oxidation ability of oxygen is higher in this case, so the reduction of Cr(VI) by Fe(II) oxidation by  $\text{O}_2$  is more likely to be strongly competed by Fe(II) and oxygen under the condition of high oxygen alkalinity. Therefore, the removal rate of Cr(VI) is lower than that in the absence of oxygen. On the other hand, in the case of hyperoxia, the reaction system continuously exchanges oxygen with the outside world, and the Fe(II) in the system is consumed more quickly, which leads to the reverse reaction of the redox reaction in the later stage, and the Cr(VI) in the system has an upward trend. However, in the anoxic group, the concentration of Cr(VI) in the solution decreased at first and then tended to be stable, because in this case, the reaction system was not affected by oxygen and other environmental factors, and the reduction effect was relatively stable.

In summary, in addition to the high pH, DO has less interference with the redox reaction of Fe(II) with Cr(VI). Therefore, Fe(II) on Cr(VI) can be divided into three stages in pH=3.0–8.0 gradient. The chemical reaction equations corresponding to each pH are summarized as follows [23]:





1. Acidic conditions (pH=3.0)



2. The hydroxide of trivalent ion and trivalent chromium is produced at pH=4.0–5.0




3. Fe(III)–Cr(III) coprecipitation is formed at pH=6.0–8.0. In the course of the experiment, it is observed that there was a red-brown flocculent precipitation in the reaction vessel



#### Effects of initial reactant concentration Fe(II):Cr(VI)

To determine whether the initial reaction concentration ratio of Fe(II) and Cr(VI) influences the form conversion and reaction rate of heavy metal pollutant Cr(VI) in the underground environment, the initial reactant concentration in the partial experiment is set as Fe(II):Cr(VI)=1:1, 2:1, 3:1, 4:1, 5:1 according to the stoichiometric ratio. To meet the actual pH environment of the ground water, the pH value is controlled to 6, and the result is shown in Fig. 5.

The results indicate that when the ratio of Fe(II) to Cr(VI) is 1:1, Fe(II) is rapidly consumed, the whole oxidation reduction reaction is incomplete, and the removal rate of Cr(VI) is relatively low with or without oxygen. When the concentration ratio of Fe(II) to Cr(VI) is 2:1, the redox system reacts violently, and there is no difference between hyperoxia and anoxic. When the concentration ratio is 3:1, the Cr(VI) was basically completely removed under the condition of anoxic. The incomplete reaction in the case of hyperoxia may be due to the fact that some Fe(II) are oxidized by oxygen and lack of effective Fe(II) to remove Cr(VI). Consequently, there is no difference between the reactant concentration ratio of 3:1 and 2:1 in the case of hyperoxia. When the concentration ratio was 4:1 and 5:1, Cr(VI) was oxidized rapidly by Fe(II) in both hyperoxia and anoxic environments, and oxygen had no obvious effect on the redox reaction of Fe(II) and Cr(VI). After 2 min, there was no residual




**Fig. 5** Variation of Fe(II) and Cr(VI) concentrations under different initial reactant concentrations

Cr(VI) concentration in the system. However, in the experimental results, the concentration of Fe(II) under the condition of high oxygen is lower than that of the oxygen-free condition, because in the presence of oxygen, the Fe(II) is oxidized by oxygen, resulting in a relatively high concentration of Fe(II) in the reaction system under anoxic conditions.

From the view of removal rate, as shown in Fig. 6, with the increase of the concentration ratio of Fe(II):Cr(VI), the removal rate of Cr(VI) in the reaction system also increased. On the whole, the removal rate of Cr(VI) in the reaction system was generally higher than that in

aerobic condition under the condition of no oxygen. Compared with other concentration, with the ratio of 3:1, the removal rate of Cr(VI) in the anoxic condition was up to 98.79%, and the reaction was basically complete. Under aerobic conditions, the removal rate of Cr(VI) was only 81.46%, which was not much different from that of 2:1 concentration ratio, which was due to the incomplete redox reaction of Fe(II) and Cr(VI) caused by competitive oxidation of Fe(II) by oxygen. When the ratio is 4:1 and 5:1, the Cr(VI) removal rate in hyperoxia and anoxic condition is more than 99%, which shows that the excess of Fe(II) and Cr(VI) in the reaction system is basically reduced, and the existence of oxygen has no obvious influence on the removal of Cr(VI), that is, the presence of oxygen is less interfering with the redox reaction of Fe(II) to Cr(VI) [27].



**Fig. 6** Removal rate of Cr(VI) at different initial reactant concentrations

**Effects of hardness and carbonate**


In actual groundwater, hardness is common, total hardness refers to the total concentration of calcium and magnesium, and carbonate hardness is a part of the total hardness, which is equivalent to the hardness formed by calcium and magnesium combined with bicarbonate in water [31]. Therefore, in the process of surface-groundwater interaction, the hardness of groundwater is one of the important factors that cannot be ignored in the morphological transformation of heavy metal Cr(VI). As shown in Fig. 7, the effects of calcium, magnesium




and bicarbonate ion on the redox reaction of Fe(II) and Cr(VI) are mainly studied in this part of the experiment, and the conditions of hyperoxia and anoxic condition are discussed, respectively.

**Effects of calcium hardness**

Combined with Fig. 8, compared with (a) and (b) figures in Fig. 7, it can be found that the removal efficiency of Cr(VI) by Fe(II) has been greatly improved after adding Ca<sup>2+</sup>. Taking hyperoxia as an example, the removal rate of Cr(VI) is 59.35% in the absence of Ca<sup>2+</sup> and 78.29% when Ca<sup>2+</sup> increases to 40 mg/L.



**Fig. 8** Removal rate of Cr(VI) in different groundwater hardness ions



**Fig. 7** Variation of Fe(II) and Cr(VI) concentrations under the influence of different groundwater hardness ions

The results describe that in the presence of Ca<sup>2+</sup> alone, the redox reaction between Fe(II) and Cr(VI) can be greatly promoted. The reason why Ca<sup>2+</sup> can promote the reaction may be that Ca(OH)<sub>2</sub> precipitation material is easily formed by Ca<sup>2+</sup> and OH<sup>-</sup> in the water and the hydroxide of the original reaction is replaced by the calcium ions to form a new ferrite coprecipitation, and the Cr(III) is further held up, resulting in a reduction in Cr(III) in the liquid phase [32], which contributes to the forward progression of the reaction in accordance with the oxidation–reduction equilibrium principle. Compared with the case without Ca<sup>2+</sup>, the concentration of Fe(II) in the reaction system also decreased to a certain extent. However, after 75 min, the concentration of Cr(VI) in the system is slightly increased, because the pH in the reaction system is stable at the pH of about 4 and the acidic environment, and the generated Ca(OH)<sub>2</sub> flocculent precipitate can be dissolved in the acid, so that the concentration of Ca(OH)<sub>2</sub> in the system is reduced, and the reaction is caused to be carried out in the reverse reaction direction, resulting in a slight increase in Cr(VI) in the system.

**Effects of magnesium hardness**

Combined with Fig. 8, compared with (a) and (c) in Fig. 7, it can be seen that the effect of Mg<sup>2+</sup> is similar to that of Ca<sup>2+</sup>. When Mg<sup>2+</sup> exists alone, it is similar to that of calcium ion, which can promote the redox reaction between Fe(II) and Cr(VI) to a certain extent. Taking hyperoxia as an example, the removal rate of Cr(VI) is 59.35% when there is no Mg<sup>2+</sup>, and the removal rate of Cr(VI) can reach 77.94% when Mg<sup>2+</sup> increases to 40 mg/L, increased by 18.59%.

**Effects of carbonate**

Compared with (a) and (d) in Fig. 7, it can be seen that the increase of HCO<sub>3</sub><sup>-</sup> in the reaction system has a

higher effect on the removal efficiency of Cr(VI) than  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ . The removal rate of Cr(VI) was 59.35% without  $\text{HCO}_3^-$ , and after adding 60 mg/L  $\text{HCO}_3^-$ , the final removal rate was 82.67%, which increased by 23.32%. This is for the reason that when  $\text{HCO}_3^-$  is added, the pH in the system changes from the original acidic environment to the alkaline environment, which reduces the Eh in the reaction system, and the redox reaction is mainly the reduction effect of Fe(II) on Cr(VI), which is beneficial to the forward reaction, so the removal rate of Cr(VI) is higher than that of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ .

**Effects of calcium hardness and carbonate**

When  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$  were added to the reaction system, the concentration was measured as  $\text{CaCO}_3$ . Compared with (a) and (e) in Fig. 7, it can be seen that the increase of  $\text{CaCO}_3$  has a very significant effect on the removal efficiency of Cr(VI).

The removal rate of Cr(VI) was 59.35% without  $\text{CaCO}_3$ , and after adding 100 mg/L  $\text{CaCO}_3$ , the final removal rate was 88.14%, which increased by 28.79%. Taking hyperoxia as an example, the reason for this phenomenon is that when  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$  are added at the same time, the  $\text{CaCO}_3$  precipitation which is insoluble in water is formed rapidly, the pH in the reaction system increases, and finally it is stable at about pH=8. The above studies show that the removal rate of Cr(VI) reaches the highest in pH=8, so the addition of  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$  at the


same time is beneficial to the redox reaction of Fe(II) and Cr(VI). Besides, the presence of  $\text{CaCO}_3$  may lead to a fraction of  $\text{Fe}(\text{OH})_3$  and  $\text{Cr}(\text{OH})_3$ , and the concentration of Fe(II) in the reaction system is smaller than that of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{HCO}_3^-$  alone, resulting in a higher removal rate of Cr(VI) than  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{HCO}_3^-$  alone.

**Effects of humic acid**



Humus is the most widely distributed natural organic matter in the earth’s ecological environment. It is considered that humus, as an important chelating agent in the environment, has a great influence on the migration and transformation of heavy metal pollutants [31]. Accordingly, humus is one of the important factors affecting Cr(VI) morphological transformation in the system of surface-groundwater interaction. Humic acid is the main component of humus. So humic acid is used to represent humus in the redox reaction of Fe(II) and Cr(VI), as is shown in Fig. 9.

**Effects of different humic acid concentration**

As can be seen from Figs. 9 and 10, the addition of humic acid (HA) can promote the redox reaction of Fe(II) and Cr(VI), and the higher the removal efficiency of Cr(VI) in the system as the HA concentration increases. In the case of hyperoxia or anoxic, the concentration of Fe(II) increased twice and then decreased when the



**Fig. 9** Changes of Fe(II) and Cr(VI) concentrations under different humic acid concentrations




HA = 15 mg/L and HA = 40 mg/L, indicating that in the redox process, HA is called the promoter of Cr(VI) reduction [33], that is to say, Fe(III), which is oxidized by Cr(VI), is converted to soluble Fe(II) again under the reduction of HA [34]. This is because Fe(III) is present in the form of a cation and Cr(VI) is present in the form of an anion, so that the affinity of Fe(III) to the HA (anion) is higher than Cr(VI) [35]. Therefore, when Fe(III) is produced by Cr(VI) reduction reaction, HA begins to take precedence over Cr(VI) and Fe(III), and the regenerated Fe(II) can continue to restore Cr(VI), to further improve the removal efficiency of Cr(VI) [33]. Even in the case of hyperoxia, humic acid has obvious reduction ability, and the soluble Fe(III) can be reduced to Fe(II), which is consistent with the results of THEIS et al. [36].

To graphically represent the mechanism of action of HA-assisted Cr(VI) morphological transformation, the conceptual model diagram can be seen in Fig. 11. In general, the mechanism can be divided into three phases: (1) when humic acid is added to the redox reaction system of Fe(II) and Cr(VI), since both Fe(II) and HA are characterized by a rapid reduction reaction at the initial phase of the reaction, Cr(VI) will first be transformed into Cr(III) under the reaction of HA and Fe(II), respectively. Fe(II) is also directly oxidized by Cr(VI) to form Fe(III). (2) When the reaction was carried out for some time, on the one hand, Fe(II) acted as the main reducing agent to reduce Cr(VI). On the other hand, the effect of HA at this time is mainly to reduce the Fe(III) produced by the oxidation of Cr(VI) to Fe(II). Because the reduction functional groups (hydroxyl and carboxyl groups) in the humic acid can complex together with the solubility Fe(III) to form a stable chelating structure, transfer electrons to the solubility

Fe(III), and the Fe(III) yields electrons to generate the solubility Fe(II) [34]. (3) As the reaction proceeded, since the reduction effect of HA on Fe(III) keeps the Fe(II) in the reaction system continuously generated, the reduction of Cr(VI) can continue. Throughout the reaction, two kinds of Fe(II) contributed: as the original Fe(II) in the phase 1 and/or 2 and the HA-reproduced Fe(II) in the phase 2 and/or 3. HA also assisted in two ways: direct reduction of Cr(VI) and reduction of Fe(III) to generated Fe(II) to continue reduction of Cr(VI).

#### Effects of humic acid on Fe(II), Fe(III), Cr(VI) and Cr(III) separately

To further determine the effect of humic acid on the parts of Fe(II), Fe(III), Cr(VI) and Cr(III), an additional experiment of the separate reaction of HA with Fe(II), Cr(VI), Fe(III) and Cr(III) was added, as shown in Fig. 12. The experimental results show that humic acid has the ability to oxidize Fe(II) alone, whether in hyperoxia or in the absence of oxygen. When only the two substances exist alone in the reaction system, HA can oxidize 42% and 52% Fe(II) to Fe(III). At the same time, humic acid has the ability to reduce Cr(VI) alone. When only these two substances exist in the reaction system, HA can reduce 25–29% of Cr(VI) to Cr(III). In addition, humic acid has the ability to reduce Fe(III) regardless of the presence of oxygen [37]. It can be found from Fig. 11 that when the Fe(III) is 12 mg/L and the humic acid is 40 mg/L, the conversion of the Fe(III) under the high oxygen condition reaches 7.76%, and the conversion rate of the Fe(III) under the anoxic condition can reach 27.89%. Furthermore, the humic acid has a certain oxidation effect on Cr(III) under the anoxic condition, and when the reaction time is between 60 and 80 min, the



concentration of Cr(VI) generated in the solution can reach 7.92 mg/L, and the conversion rate is 66%.

The above results can explain that when humic acid, Fe(II) and Cr(VI) exist together, that is, when the reaction time is 60 to 80 min, the concentration of Cr(VI) in the reaction system increases briefly. However, in the whole redox process, humic acid is still dominated by reduction, so the generated Cr(VI) will be immediately reduced to Cr(III).

## Conclusions

In the present investigation, the interaction process of surface-groundwater was simulated by changing the dissolved oxygen content in the redox reaction system of dissolved Fe(II) and Cr(VI). The main conclusions are as follows:

The content of DO does affect the redox of Fe(II) on Cr(VI). Compared with anoxic condition, it is more favorable for Fe(II) to fix Cr(VI) under aerobic condition,

that is, during the contact of ground water and surface water, the change of DO concentration in the reaction system is beneficial to the conversion of the heavy metal pollutant Cr(VI) to the low-valence Cr(III) by the underground mineral element Fe(II).

pH has a great influence on the redox reaction of Fe(II) with Cr(VI). The reaction rate of Fe(II) with Cr(VI) and the removal rate of Cr(VI) decrease with the increase of pH at pH = 3–5, and increase with the increase of pH at pH = 6–8. When it comes to the condition of surface-groundwater interaction, that is, when the concentration of DO changes, the Fe(II) is superior to O<sub>2</sub> and Cr(VI) in redox reaction when the pH = 3–5; when the pH value is 6 to 8, the Fe(II) is more easily oxidized by the O<sub>2</sub> under the condition of hyperoxia and alkali. The reduction of Cr(VI) by Fe(II) is strongly competed by the reaction of Fe(II) with oxygen. The conversion of Cr(VI) in the case of high oxygen is lower than that in the absence of oxygen.

The effect of DO on the redox reaction of Fe(II) and Cr(VI) is closely related to the initial concentration ratio of Fe(II) to Cr(VI). When Fe(II):Cr(VI) is 1:1, 2:1, 4:1 and 5:1, DO has no obvious effect on the reaction rate and Cr(VI) removal rate. When the Fe(II):Cr(VI) is 3:1, the removal rate of Cr(VI) is 98.79%, which is a complete reaction, under the condition of anoxic. The influence of DO on the reaction results in the partial Fe(II) being oxidized by the oxygen and the absence of sufficient and effective Fe(II) to convert the Cr(VI) with hyperoxia condition.

In the presence of groundwater hardness ions Ca<sup>2+</sup>, Mg<sup>2+</sup> and HCO<sub>3</sub><sup>-</sup> alone, the redox reaction between Fe(II) and Cr(VI) is greatly promoted, which is beneficial to the conversion of Cr(VI) to Cr(III). When Ca<sup>2+</sup> and HCO<sub>3</sub><sup>-</sup> exist at the same time, they can promote the original reaction system more. Therefore, increasing the hardness of groundwater is beneficial to the conversion of toxic Cr(VI) into nontoxic Cr(III).

When there is HA, Fe(III) formed by the oxidation of Fe(II) by Cr(VI) will be converted to soluble Fe(II) again under the reduction of HA. Therefore, HA can promote the redox reaction between Fe(II) and Cr(VI), and promote the conversion of Cr(VI) by Fe(II) more effectively. With the increase of HA concentration, the removal rate of Cr(VI) in the system is higher.

Combined with the above results, in the process of sub-surface-groundwater interaction with changing actual oxygen content, to achieve a better transformation/removal effect of Cr(VI) in the interaction band, the following measures may be taken: appropriate adjustment of pH to neutral or weak alkalinity, adding external soluble Fe(II) reagent, increasing hardness of groundwater and humic acid concentration. In these experimental conditions, such as increasing Fe(II):Cr(VI) to above 3:1,

adding 60 mg/L  $\text{HCO}_3^-$ , adding 100 mg/L  $\text{CaCO}_3$ , adding 40 mg/L HA, etc., we can achieve Cr(VI) removal rate of more than 80% or even more than 99%.

#### Abbreviations

DO: Dissolved oxygen; Cr(VI): Hexavalent chromium; Fe(II): Ferrous iron; Cr(III): Trivalent chromium; Fe(III): Ferric iron; DI: Deionized water;  $\text{K}_2\text{CrO}_4$ : Potassium bichromate;  $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2$ : Ammonium ferrous sulfate;  $\text{HNO}_3$ : Hydrogen nitrate; NaOH: Sodium hydrate;  $\text{CaCl}_2$ : Calcium chloride;  $\text{MgCl}_2$ : Magnesium chloride;  $\text{NaHCO}_3$ : Sodium hydrogen carbonate; HA: Humic acid.

#### Acknowledgements

We are highly grateful to Key Laboratory of Yangtze River Water Environment for provision of testing instruments, and to School of Environment and Geographical Science for provision of laboratories.

#### Authors' contributions

Concept of the study, experiment execution and manuscript writing: YZ; experimental guidance and manuscript revision: CD, SL, YD. All authors read and approved the final manuscript.

#### Funding

This study was funded by the National Natural Science Foundation of China (41672230, 41601514, 51961145106), Shanghai "Science and Technology Innovation Action Plan" Project (19230742400, 19ZR1459300), Shanghai Peak Discipline Project (0200121005/053, 2019010202), State Key Laboratory of Petroleum Pollution Control (PPC2016019).

#### Availability of data and materials

All data generated or analyzed during this study are included in this article.

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declared that they have no competing interests.

#### Author details

<sup>1</sup> Department of Hydraulic Engineering, Tongji University, Shanghai, China. <sup>2</sup> Key Laboratory of Yangtze River Water Environment, Ministry of Education, Shanghai, China. <sup>3</sup> School of Environmental and Geographical Sciences, Shanghai Normal University, Shanghai, China. <sup>4</sup> Institute of Water and Ecology Problems, Far East Branch of the Russian Academy of Sciences, Khabarovsk, Russia. <sup>5</sup> Department of Environmental Engineering, University Tunku Abdul Rahman, Perak, Malaysia. <sup>6</sup> Department of Civil and Environmental Engineering, Pontifical Catholic University of Rio de Janeiro, Rio de Janeiro, Brazil.

Received: 9 January 2020 Accepted: 23 March 2020

Published online: 15 April 2020

#### References

- Andaleeb F, Zia MA, Ashraf M, Khalid ZM (2008) Effect of chromium on growth attributes in sunflower (*Helianthus annuus* L.). *J Environ Sci* 20:1475–1480
- Tomaszewski EJ, Lee S, Rudolph J, Xu H, Ginder-Vogel M (2017) The reactivity of Fe(II) associated with goethite formed during short redox cycles toward Cr(VI) reduction under oxic conditions. *Chem Geol* 464:101–109
- Hug SJ, Buerge IJ, Weidler PG (1997) Transformations of chromium in the environment. *Anal Mag* 25:12–15
- Haddad HH (2012) The effect of heavy metals cadmium, chromium and iron accumulation in human eyes. *Am J Anal Chem* 3:710–713
- Blasiak J, Kowalik J (2000) A comparison of the in vitro genotoxicity of tri- and hexavalent chromium. *Mutat Res Genet Toxicol Environ Mutagen* 469(1):135–145
- Datta S, Mailloux B, Jung HB et al (2009) Redox trapping of arsenic during groundwater discharge in sediments from the Meghna riverbank in Bangladesh. *Proc Natl Acad Sci USA* 106:16930–16935
- Huili Du (2007) Discussing on groundwater flow near partially penetrating river. Hohai University, Nanjing
- Harvey JW, Fuller CC (1998) Effect of enhanced manganese oxidation in the hyporheic zone on basin-scale geochemical mass balance. *Water Resour Res* 34:623–636
- Lasagna M, Luca DAD, Franchino E (2016) Nitrate contamination of groundwater in the western Po Plain (Italy): the effects of groundwater and surface water interactions. *Environ Earth Sci* 75(3):1–16
- Mclachlan PJ, Chambers JE, Uhlemann SS, Binley A (2017) Geophysical characterization of the groundwater–surface water interface. *Adv Water Resour* 109:302–319
- Schwaezenbach RP, Giger W, Hoehn E et al (1983) Behavior of organic compounds during infiltration of river water to groundwater. *Field studies. Environ Sci Technol* 17(8):472–479
- Chaubey J, Arora H (2017) Transport of contaminants during groundwater surface water interaction. In: *Development of water resources in India*, pp 153–165
- Woessner WW (2010) Stream and fluvial plain groundwater interaction-rescaling hydrologic thought. *Ground Water* 38(3):423–429
- Winter TC (1999) Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrogeol J* 7(1):28–45
- Sophocleous M (2010) Managing water resources systems: why "Safe Yield" is not sustainable. *Ground Water* 35(4):561
- Xue Y, Zhu X, Wu J, et al (1997) *Underground water dynamics*. Geological Press
- Charette MA, Sholkovitz ER, Hansell CM (2005) Trace element cycling in a subtropical estuary: part 1. Geochemistry of the permeable sediments. *Geochim Cosmochim Acta* 69:2095–2109
- Wu P, Jin J, Sun D, et al (2011) Hydrogeochemical study of surface water-ground water interactions in Xinanquan domain. 1st edn, China Water Conservancy and Hydropower Press
- Gambrell RP (1994) Trace and toxic metals in wetlands—a review. *J Environ Qual* 23(5):883–891
- Singh IB, Singh DR (2002) Influence of dissolved oxygen on aqueous Cr(VI) removal by ferrous ion. *Environ Technol Lett* 23(12):1347–1353
- Tang S, Wang X, Mao Y et al (2015) Effect of dissolved oxygen concentration on iron efficiency: removal of three chloroacetic acids. *Water Res* 73:342–352
- Shiqiang W, Guanglin L, Sterberg R, Dingyong W (2003) Kinetics of metal ion reactions with humic acids and index for steady-state. *Acta Pedol Sin* 31(1):132–134
- Buerge IJ, Hug SJ (1997) Kinetics and pH dependence of chromium (VI) reduction by iron (II). *Environ Sci Technol* 31:1426–1432
- Eary LE, Rai D (1988) Chromate removal from aqueous wastes by reduction with ferrous ion. *Environ Sci Technol* 22(8):972–977
- Binyuan W et al (2014) Effect of pH value on reduction treatment of chromium-containing wastewater by ferrous and kinetics study. *J Nat Sci Heilongjiang Univ* 31(3):356–360
- Sedlak DL, Chan PG (1997) Reduction of hexavalent chromium by ferrous iron. *Geochim Cosmochim Acta* 61:2185–2192
- Schlautman MA, Han I (2001) Effects of pH and dissolved oxygen on the reduction of hexavalent chromium by dissolved ferrous iron in poorly buffered aqueous systems. *Water Res* 35(6):1534–1546
- Yoon I-H, Bang S, Chang J-S et al (2011) Effects of pH and dissolved oxygen on Cr(VI) removal in Fe(0)/ $\text{H}_2\text{O}$  systems. *J Hazard Mater* 186:855–862
- Stumm W, Morgan JJ (1996) *Aquatic chemistry*, 3rd edn. Wiley Interscience, New York
- Salem FY, Parkerton TF, Lewis RV et al (1989) Kinetics of chromium transformations in the environment. *Sci Total Environ* 86(1–2):25–41
- Zhou M (2007) The effect of humic acids on Zero Value Iron (ZVI) reduction of Cr(VI) in contaminated water. Zhejiang University



32. Lo IMC, Lam CSC, Lai KCK (2006) Hardness and carbonate effects on the reactivity of zero-valent iron for Cr(VI) removal. *Water Res* 40(3):595–605
33. Hori M, Shozugawa K, Matsuo M (2015) Reduction process of Cr(VI) by Fe(II) and humic acid analyzed using high time resolution XAFS analysis. *J Hazard Mater* 285:140–147
34. Li J, Luan F, Xie L et al (2009) Study on the influencing factors of humic acid reduction of Fe(III). *Environ Pollut Control* 31(2):23–30
35. Zhilin DM, Schmitt-Kopplin P, Perminova IV (2004) Reduction of Cr(VI) by peat and coal humic substances. *Environ Chem Lett* 2:141–145
36. Theis TL, Singer PC (1974) Complexation of iron (II) by organic matter and its effect on iron (II) oxygenation. *Environ Sci Technol* 8(6):569–573
37. Wang Q, Wei S (2005) Studies on the dynamics of oxidation–reduction of Fe<sup>3+</sup> with humic acids and fulvic acids. *China Environ Sci* 25(4):480–483
38. Tong M (2015) Activation of O<sub>2</sub> by Fe(II) in subsurface environments for reactive oxygen species production and arsenic removal. China University of Geosciences, Wuhan

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

---